Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.





United States Department of Agriculture
Science and Education Administration, Agricultural Research
North Central Region



Source:

Spring and Durum Wheat Quality Laboratory USDA, SEA, Agricultural Research Cereal Chemistry & Technology, N.D.S.U. Fargo, North Dakota 58105



REPORT OF PHYSICAL, CHEMICAL, MILLING AND BAKING EXPERIMENTS WITH HARD RED SPRING WHEAT

1980 CROP1/


by

R. D. Maneval, Food Technologist; R. D. Crawford, A. A. Ottenbacher, Technicians, D. W. Lillard, Res. Chemist, J. E. Wohlman, Secretary, SEA/Agricultural Research; 2/ L. L. Nolte, and M. Skunberg, Technicians, NDSU; 3/ and V. L. Youngs, Research Leader.2/

This is a progress report of cooperative investigations containing some results that have not been sufficiently confirmed to justify general release; interpretations may be modified with additional experimentation. Confirmed results will be published through established channels. Cooperators submitting samples for analysis have been given analytical data on their samples prior to release of this report. The report is primarily a tool for use of cooperators and their official staffs and to those persons having direct and special interest in the development of agricultural research programs.

This report was compiled by the Science and Education Administration, Agricultural Research, U. S. Department of Agriculture. Special acknowledgment is made to the North Dakota State University for their facilities and services provided in support of these studies. The report is not intended for publication and should not be referred to in literature citations nor quoted in publicity or advertising. Use of the data may be granted for certain purposes upon written request to the agency or agencies involved.

^{2/} Hard Red Spring & Durum Wheat Quality Lab., NDSU.
3/ Dept. of Cereal Chemistry & Technology, NDSU.



1980 COOPERATING AGENCIES, STATIONS AND PERSONNEL

The cooperative agencies and stations conducting the varietal plot and nursery experiments from which the 1980 spring wheat samples were received are listed below:

Arizona Agricultural Experiment Station:
Mesa

Idaho Agricultural Experiment Station:
Aberdeen and Tetonia

Minnesota Agricultural Experiment Station:
Crookston, Morris and St. Paul

Montana Agricultural Experiment Station:
Bozeman, Havre and Sidney

North Dakota Agricultural Experiment Station:

Carrington, Dickinson, Langdon, Minot and Williston

South Dakota Agricultural Experiment Station:
Brookings, Redfield and Selby

Washington Agricultural Experiment Station:
Pullman

Wisconsin Agricultural Experiment Station:
Madison

Wyoming Agricultural Experiment Station:
Sheridan and Torrington

A complete list of all cooperating agencies, stations, and personnel for the year will be found in the report by R. H. Busch, et al., Wheat Varieties Grown in Cooperative Plot and Nursery Experiments in the Spring Wheat Region in 1980.



INTRODUCTION


Samples of standard varieties and many of the new strains of hard red spring wheat grown in cooperative experiments in the spring wheat region of the United States 4/ have been milled each year by the USDA. The flours were assayed chemically and physically and baked into bread to determine the quality characteristics. The purpose of this report is to make available to the cooperators, quality data on the standard varieties and new strains of hard red spring wheat from the 1980 crop.

The same general format and techniques were used in evaluating the wheat as outlined in quality reports for previous years. The data contained in this report are comparable to data in past reports and, where applicable, average results and also the average results of other crop years are compared. The area averages are tabulated for the Uniform Regional Nursery varieties of Butte, Era, Chris and Waldron. A four-year average (4-YA) and the averages for the individual four years include all selections grown in the Uniform Regional Nurseries for that year. These results give an overview of individual years and the influence of environment on the crop. The actual crop characteristics may be somewhat different due to differences in varieties, but the change from year to year is applicable.

The evaluation of a sample involves three areas of analysis: kernel characteristics, milling performance and baking evaluation. A brief description of the technique is given on pages 11 and 15 of this report. It is possible to quickly deduce the various characteristics of the selection and any outstanding features or deficiencies which are apparent. No specific comments are made regarding the mixogram patterns, since reference mixograms for each of the general types are presented at the end of the report.

Seeding for the 1980 crop over the spring wheat area was early to normal with severe drought conditions prevailing over much of the area until August. Some areas experienced excess rainfall during harvest which caused some problems. The average flour extraction was 0.3% higher than the 1979 crop and 0.3% lower than the 4-YA.

^{4/} Busch, R. H., and Quick, J. S. Wheat Varieties Grown in Cooperative Plot and Nursery Experiments in the Spring Wheat Region in 1980. SEA/Agricultural Research, U. S. Department of Agriculture and State Agricultural Experiements Station, St. Paul, MN.



Wheat mineral content was slightly higher than the 1979 crop and the 4-YA. The wheat protein concentration was 0.7% higher than the 1979 crop average and 0.1% higher than the 4-YA. The physical characteristics of the wheat were down slightly from the 4-YA.

The baking performance of the 1980 crop was slightly stronger than the 1979 crop. Mixing times were longer and the dough properties were slightly stronger. Absorption was 1.6% higher than the 4-YA. Flour protein was 0.3% higher than the 4-YA and 1.0% higher than the 1979 crop. Loaf volume was also slightly better than the 4-YA. Oxidation requirements were unchanged.



SOURCE OF THE 1980 CROP SAMPLES

Tests were performed on 1,204 samples. However, data on 527 of these are not included in this report, because this information was of interest to plant breeders at specific experiment stations only. Data presented in this report are from the Field Plot Nursery, Uniform Regional Nursery, International Sawfly Yield Nursery and the International Spring Wheat Nursery. The samples came from the 21 stations in nine states shown below:

Arizona: Mesa

Idaho: Aberdeen and Tetonia

Minnesota: Crookston, Morris and St. Paul

Montana: Bozeman, Havre and Sidney

North Dakota: Carrington, Dickinson, Langdon,

Minot and Williston

South Dakota: Brookings, Redfield and Selby

Washington: Pullman Wisconsin: Madison


Wyoming: Sheridan and Torrington

On page 7 are listed the spring wheats that were included in the Uniform Regional Nursery trials. The variety or cross, the station that developed the variety, the state selection number and the C.I. number are given.

BLENDING AND AVERAGING PROCEDURES USED

Individual wheat samples from the Uniform Regional Nursery originating from the three geographical areas shown in the illustration on page 6 were blended according to area. Only 18 of the 21 stations were compatible for blending. Milling performance, mixograms and baking data were obtained from these area blends. However, data for kernel characteristics are arithmetical averages of individual sample analyses. These data from the Uniform Regional Nursery also are compared with averages from the previous four years (Table 4).





Wheat blends were made according to the geographical areas shown above.

Data from the International Sawfly Yield Nursery were obtained on the individual samples. Table 14 gives the variety averages for the 1980 crop year.

Data for the Field Plot Nursery and the International Nursery are on the individual samples.




ENTRIES IN THE 1980 UNIFORM REGIONAL

HARD RED SPRING WHEAT NURSERY

En hass	Cross or	OT No. 27	37	
Entry	Cross or	CI No. or	Year	Course
No.	Variety	Sel. No.	Entered	Source
1.	Marquis	3651	1929	Canada
2.	Chris	13751	1969	USDA-MN
3.	Waldron	13958	1964	ND
4.	Era	13986**	1972	USDA-MN
5.	Butte	17681	1979	ND
6.	Olaf/MN6792	SD2700	1980	SD
7.	Butte/Eureka	SD2835	1980	SD
8.	Hand/2*WS1809/3/	SD2870	1980	SD
	Oleson/Tobari//Trapper			
9.	Butte/James 's'	SD2868	1980	SD
10.	ND476/4/Sheridan/3/	MT648*	1979	USDA-MT
	Norin 10/Bev. 14//4*			
	Centan			
11.	TzPP/Son 64//Selkirk	MT7836**	1980	USDA-MT
12.	Mrn/Tbr 66/3/TzPP/	ID0167**	1980	USDA-ID
	AN3//B61-136			
13.	Neepawa*6/RL4137	RL4352	1980	MantCanada
14.	Waldron/Era	MN70170R**	1980	USDA-MN
15.	Crim/Era*2//Bui-Gallo		1979	USDA-MN
16.	Crim/Era*2//MN6923S	MN7357**	1980	USDA-MN
17.	Era*/Chris M	MN7222**	1977	USDA-MN
18.	Olaf//ND499/ND516	ND569**	1979	ND
19.	Butte*2/ND507	ND572	1980	ND
20.	ND526/Kitt	ND573	1980	ND
21.	Olaf/5/Jt/ND335/	ND574**	1980	ND
	Pembina/3/Wanken 2/4/			
	Cis/Wisc 261	370 F 7 F	1000	ND
22.	ND551//Butte*2/ND507	ND575 NK75S2634**	1980 1980	Northrup King
23.	Waldron/Era	ND75S2631**	1980	Northrup King
24.	Era/Justin	HS7664**	1980	No. Amer. Pl. Br.
25.		HS74183**	1978	No. Amer. Pl. Br.
26.		X6753**	1980	Pioneer
27.		X6718**	1980	Pioneer
28.	N70000E2/E710E2	WA6756**	1980	WA
29. 30.	N7000052/K71053 K71051/WA5949	WA6758**	1980	WA
30.	K / TOJI/ WAJJ47	1120750		

^{**} Semidwarf



METHODS

The terminology and methods used are briefly described below:

Test Weight Per Bushel - The weight per Winchester bushel of cleaned, dry, scoured wheat. To determine the dockage-free test weight on a comparable sample, approximately one pound per bushel should be subtracted from the value given.

1000 Kernel Weight - The 1000 kernel weight was determined by counting with a Seedburo seed counter the number of kernels in a 10 g sample of cleaned, picked wheat5/.

Kernel Size - The percentages of the size of the kernels (large, medium and small) were determined on a wheat sizer as described by Shuey6/.

The sieves of the sizer were clothed as follows:


Top Sieve - Tyler #7 with 2.92 mm opening Middle Sieve - Tyler #9 with 2.24 mm opening Bottom Sieve - Tyler #12 with 1.65 mm opening

Potential Yield - The potential yield is not shown on the computer tables, but it can be determined by multiplying the percentages of the overs of each sieve #7, #9 and #12 by the value of 78%, 73% and 68%, respectively. The accumulation percentage would be the potential yield.

Milling - The samples were cleaned by passing the wheat over an Emerson kicker and dockage tester and through a modified Forster scourer (Model 6). The clean, dry samples were pretempered to 12% moisture for at least 72 hours; then tempered to 16% moisture and allowed to stand overnight prior to milling.

^{5/} Mention of a trademark name or a proprietary product does not constitute a guarantee or warranty of the product by the U. S. Department of Agriculture, and does not imply its approval to the exclusion of other products that may also be suitable.

^{6/} Shuey, William C. A Wheat Sizing Technique for Predicting Flour Milling Yield. Cereal Science Today 5:71-72,75 (1960).



The International Sawfly Nursery, Special Uniform Nursery and the International Spring Wheat Yield Nursery samples were milled on a Brabender Quadrumat Jr. mill. The mill was equipped with a #18 wire on the drum sieve. The throughs of the #18 wire were rebolted on a Strand sifter equipped with a #60 Tyler sieve. The sample was sifted for 1 minute. The throughs of the #60 wire were classified as flour, and this was the material tested. The overs of the #18 wire were classified as bran, and the overs of the #60 Tyler sieve as crude shorts.

The Uniform Regional Nursery blends and the Field Plot Nursery samples were milled on a Buhler continuous experimental mill. This mill has been slightly modified to give results more comparable to commercial milling. The break scalping sieves were clothed with #54 stainless steel wire, the reduction scalping sieves with #58, #66 and #105 stainless steel wire for the first, second and third reduction, respectively. All of the flour sieves were clothed with #135 stainless steel wire.


All six flour streams were combined to give the patent flour. The extraction of a good milling wheat using this flow is approximately 68%. This is comparable to a commercial "long patent" extraction flour. At this flour extraction of the wheat, the changes in flour ash are most sensitive to changes in percent extraction.

<u>Protein Content</u> - The protein was calculated by multiplying the factor of 5.7 times the percent nitrogen as determined by the standard Kjeldahl procedure.

Mineral Content or Ash Content - This was determined by measuring the residue of the minerals left after incinerating the sample for approximately 16 hours at 565°C. The results were reported as percentage of the sample that was incinerated.

Mixogram - The mixogram was determined by using 30 g of flour and adding 20 cc of water. The sensitivity spring setting was set at 10. All mixograms were run with constant weight of flour and volume of water. Absorptions reported were adjusted according to the height of the mixogram. The correction factor was determined from a series of flours by varying the amount of absorption.

Mixogram Pattern - The reference mixogram patterns given at the end of the report demonstrate the different types of mixograms that were obtained. A single number is assigned each pattern to characterize and simplify the classification of the curves--the larger number indicating stronger curve characteristics.



Baking Procedure or Formula - The baking formula used was as follows:

100% flour 3% milk D.S.M.

2% salt 3% yeast

5% sugar 2% shortening (Crisco, melted)


The samples were mixed to development in National Manufacturing mixers: the macro mixer for the 25 g samples and the 100 g special mixer for the 100 g samples. Bromate (7.5 ppm) for oxidation and barley malt flour (0.1%) for enzymatic supplement were added to each sample. All doughs were moulded in a Roll-Er-Up moulder.

Absorption - The amount of water, expressed as percent of the flour, required to bring the dough to proper consistency.

Crumb Color - A value was determined by comparing the loaf of the tested sample against a baking standard. This standard was selected as an average for the crop year for the spring wheat area.

Loaf Volume - The volume of the baked loaf as determined by seed displacement.

All values (protein, ash and absorption) were reported on a 14% moisture basis.




DISCUSSION

The following discussion presents some of the basic techniques and criteria used in the milling and baking quality evaluation of the samples. There are four major evaluation categories used: kernel characteristics, to characterize the kernel; milling performance, to evaluate the general milling characteristics; mixogram patterns, to classify the flour as to type; and baking evaluation, to rate the flour as to overall baking.

Each evaluation category can be important. A sample could be of a sufficiently poor quality for a given category to suggest elimination from future testing. However, a sample submitted for the first time and found to be questionable should be tested again to establish if it has a satisfactory or unsatisfactory classification. A sample which is consistently rated as questionable should be discarded.

Six characteristics (test weight, 1000 kernel weight, percent large kernels, percent small kernels, wheat mineral and wheat protein) were independent variables used to calculate the dependent variable - Kernel Characteristics. Four characteristics (percent extraction, mineral @65% extraction, milling characteristic, and protein difference between flour and wheat protein) were used to calculate the dependent variable - Milling Performance. Bake absorption, mixing time, dough characteristics, crumb color, crumb grain and loaf volume were the six independent variables used to determine the dependent variable - Baking Evaluation. These three dependent variables after calculation become independent variables used to calculate the dependent variable - General Evaluation.


The three dependent variables, Kernel Characteristics, Milling Performance and Baking Evaluation are rated on a scale of 1 to 8, with 1 being Very Satisfactory and 8 being Unsatisfactory. The General Evaluation is rated on a scale of 1 to 4, with 1 being no promise; 2, little promise; 3, some promise; and 4, good promise. If one of the independent variable's converted value is 8 (with the exception of crumb color), this automatically will rate the General Evaluation as 1, or no promise. If there are no 8's, the three values are employed in a regression equation to derive the General Evaluation. The weighted value for each of these variables on the General Evaluation is approximately 6% for Kernel Characteristics, 47% for Milling Performance and 47% for Baking Evaluation.



To quickly point out problem areas for a selection, two additional columns appear on the printout. One column is Minor Deficiencies in which the independent variables, converted to a 5 or a 6 (i.e., Questionable or Questionable to Unsatisfactory) will appear. The second column is Major Deficiencies in which the independent variables were converted either to a 7 or an 8 (i.e. Unsatisfactory to Questionable and Unsatisfactory). Deficiencies of the various selections may be readily determined by scanning these columns. It is also possible to have one or two independent variables that would appear in the major deficiency These characteristics should be given column, rating 7. serious consideration even though they do not influence the general rating sufficiently to rank the selection as having no promise.

All samples, as in previous years, are compared with a milling and baking standard that represents a blend of the crop year blended to a known quality. However, the samples for the individual stations are evaluated against the average results of the check varieties from the respective The agronomic and climatic conditions of the stations. individual locations can affect the quality of the wheat sample, such that the evaluation at certain locations could have all samples--even the named varieties--classified as Questionable to Unsatisfactory. Therefore, the evaluation ratings of one station are not directly comparable with those of another station. For example, an area may produce low protein wheats which give large and plump kernels, good milling and kernel characteristics, but low protein and unsatisfactory baking properties such as short mixing time, low loaf volume and weak dough characteristics. The wheat from this area could not be considered as a strong spring wheat and would not maintain the quality expected from the spring wheat producing area. A good variety should have tolerance to a wide range of environmental conditions and the overall picture should be taken into consideration for establishing these varieties.

Kernel Characteristics are important in determining the initial value of the wheat and, if extremely poor, could disqualify a new variety from further consideration. Because of the present grading system, it is desirable to have a good test weight. If a sample has a low 1000 kernel weight and small kernel size distribution, it would be considered a poor sample for milling because of the high ratio of bran to endosperm. Therefore, it is desirable to have plump kernels. Wheat ash is an important factor when comparing a variety against other standard varieties. If a sample consistently has higher wheat mineral content, it increases the probability of having high flour ash. Lower




protein than the standard varieties is not desirable, because in a low protein crop year the probability of it having such a low protein as to be undesirable is much greater. Therefore, the protein must also be considered as a characteristic when comparing varieties grown in the same locality.

Milling Performance is very important, especially the subcategory of milling characteristics. If low extraction or high flour ash is obtained, these become major factors which are quite unacceptable from a commercial milling standpoint. All flour mineral contents are reported at a constant extraction of 65%, so that the figures are directly comparable. As a rule of thumb, one can approximate that each point of ash (0.01%) is equivalent to approximately 2% in extraction.

Milling characteristics are important. A sample which tends to be soft in character requires a different milling technique to be milled properly. On commercial mills flowed for hard vitreous spring wheats, soft milling characteristics cause great difficulty. Therefore, if a sample shows softness in character, it is considered to be unsatisfactory. Likewise, a sample which is extremely hard and vitreous will cause difficulty. Both types of wheat (soft and vitreous) require different roll pressures, clothing, sifter surface and temper to be milled properly. If these wheats are blended with normal milling wheats, improper results are obtained since these characteristics are not necessarily compatible or additive. Normal to soft score indicates that the sample shows a tendency toward softness of character on the flour mill stocks and extraction. would indicate that the sample may give some difficulty for certain mill streams, and an adjustment would either have to be made in the milling flow or in tempering procedures to compensate for these differences. The properties of this wheat may or may not be compatible with other wheats with which it may be blended; therefore, it is important to maintain varieties with milling characteristics as uniform as possible.

The amount of protein recovered in the flour for a sample is of importance. High protein wheats yielding low protein flours are not desirable. Such a wheat would have much of the protein distributed in the outer portion of the kernel which would result in excessive protein in the feed. Therefore, higher wheat protein would be necessary to yield a flour with protein content comparable to that of a wheat that gives good flour protein recovery.



Mixogram Patterns and Farinogram Patterns are important in estimating the strength and mixing tolerance or potential mixing tolerance of a flour. A long, flat curve is more desirable that a short, peaked curve; however, an extremely long curve may be undesirable, if the flour would require excessive mixing for proper development. Both the pattern and length of the curve are important, and both must be considered. Abnormal curves, such as sway-back or long initial time to incorporate the water, indicate undesirable characteristics.


Baking Evaluation takes into account the flour absorption, mixing time, dough characteristics, loaf volume and machinability. A sample which has low absorption would be unsatisfactory. A sample with extremely short mixing time would also be considered undesirable as a good strong spring wheat. When a sample is in the minimal range for these values, it is considered to be questionable until further testing demonstrates whether a definite deficiency exists.

Doughs having mellow to weak dough properties show a tendency towards weakness. Also, for mellow to strong, the dough is mellow but has a tendency to be strong, and a strong to mellow dough is just the reverse. Since these characteristics are subjective rather than objective, it is necessary at times to estimate the tendency; therefore, the necessity exists for apparent double grades.

The grain or appearance of the interior of the loaf shows how well the sample stood up during baking and may point out or explain some deficiencies which have been observed during the baking test.

Loaf volume indicates potential strength of the flour in a different manner than mixing time or dough characteristics in that it shows the ability or lack thereof for the dough to expand under pressure and to contain the entrapped gases during this expansion. Weak flours act much like rotten balloons, which burst when blown up and collapse and yield low loaf volume or extremely large volume and large holes in the interior of the loaf. Low protein flours and lifeless (dead) doughs exhibit properties similar to putty and do not expand during fermentation or baking and give low loaf volume. Tough and very bucky doughs are bound too tightly and impede expansion of the gases causing low loaf volume.

General Evaluation rating applies only to the data contained in the year of the report. A new category, the Prospect of a selection, will apply to two or more years of data. The Prospect is given for each selection that has been tested for at least two crop years. This evaluation



takes into account the various grading factors and the results of the crop years in an effort to determine if the selection should be considered as a prospective new variety. The main defects and outstanding features are discussed. A selection which is promising should be continued. Those which show some promise with outstanding agronomic characteristics should be seriously considered and looked at in large plots (if it has not been done previously), providing sufficient other information has been obtained. A sample which shows little or no promise should be discontinued.



UNIFORM REGIONAL NURSERY SAMPLES - 1980 CROP

Discussion of Area Blends


A total of 572 Uniform Regional Nursery samples were received. The samples were from 19 stations in eight states. Wheat blends were made of the samples for this crop year by area. The areas tend to represent movement of the wheat in the market. Kernel characteristics were determined on individual samples to eliminate possible erroneous results. The area blends were then milled and baked by our macro method. Thirty samples were received from each of the 19 stations. Twenty-five selections were included for quality evaluation in the Uniform Regional Nursery samples. The remainder of the samples were commercially named varieties, namely, Butte, Chris, Era, Marquis and Waldron.

Data from the northeast area blend are given in Table 1. The four stations included in this blend were Carrington, Langdon and Minot, North Dakota and Crookston, Minnesota.

- The data for the southeast area blend are given in Table 2. The five stations included in this blend were: Morris and St. Paul, Minnesota; Brookings, Redfield and Selby, South Dakota. Madison, Wisconsin was not included in the area blend because it showed sprout damage; however, the station was milled and baked separately, and the data is reported later in Table 7.
- The data for the western area blend are given in Table 3. The seven stations included in this blend were: Aberdeen, Idaho; Havre and Sidney, Montana; Dickinson and Williston, North Dakota; and Sheridan and Torrington, Wyoming. Tetonia, Idaho and Pullman, Washington were not included in the area blend because samples from Tetonia showed frost damage, and the samples from Pullman were not compatible (low protein) with the other samples from that area. Samples from these two stations were processed individually, and the data are reported in Tables 5 (Tetonia) and 6 (Pullman).

Discussion of Area and Crop Year Averages

In Table 4 are given the average area results for the combined data of the varieties, Butte, Chris, Era and Waldron samples submitted from the six states and sixteen stations. The area average represents all samples that were grown in that area for the year cited.




The milling and baking results were obtained from the area blend of the wheats in equal proportions from each of the stations for the respective variety or selection. The regular 100 g straight dough rich formula baking procedure was used in baking. The General Evaluation column includes the overall performance of the blend of each sample. The General Evaluation given for the sample area blend may not agree with that of the individual wheat samples within the blend, since averages do not express the range, and poor characteristics may be masked. In an endeavor to clarify this problem, the average general evaluation, the number of total deficiencies and the number of major deficiencies are shown in parenthesis after each variety or selection — (Average General Evaluation — #Total Deficiencies/#Major Deficiencies).

For simplicity and brevity of the report, as in previous reports, each selection or variety will be discussed from the general viewpoint rather than the individual areas. General Evaluation summarizes the results from the individual areas for one crop year. The evaluation is more meaningful for the overall performance of a variety or selection, when at least two or more crop years are included. Data discussed under the category, The Prospect, includes two or more years.

Also given in Table 4 are comparisons of the previous four crop years, which include all selections grown in the Uniform Regional Nursery for that year, as well as the 4-YA. The 1980 crop kernel characteristics (test weight, 1000 kernel weight and wheat protein) were similar to the 4-YA. Milling extraction was down slightly from the 4-YA; however, flour protein was 0.3% better than the 4-YA. Bake absorption was 1.6% higher and mixing time one-half minute longer than the 4-YA. Dough characteristics were slightly stronger and crumb color, crumb grain and loaf volume were similar to the 4-YA.

A comparison of the 1980 and 1979 crop results showed the 1980 crop to be slightly better. In general, the kernel characteristics (1000 kernel weight and test weight) were similar, with wheat protein 0.7% higher than the 1979 crop. Milling was quite similar to the 1979 crop; however, flour protein was 1.0% better than the 1979 crop. The 1980 baking absorption was 2.4% higher than in 1979 with one-half minute longer mixing time and a dough similar to the 1979 crop. Crumb color, crumb grain and loaf volume were also similar to the 1979 crop.



Discussion of Individual Varieties or Selections

Average results of the varieties Butte, Chris and Waldron for each of the individual areas were used as standards for the other selections from that area; therefore, a variety or selection may be rated satisfactory in two different areas, but comparison of the data may show much poorer results for one area due to adverse environmental conditions. Thus the sample with poor results could be rated as having unsatisfactory quality when compared with the overall spring wheat area, even though it may be rated as showing good promise for one area.

By using the same format as used in previous years and employment of the computer, all named varieties receive a general evaluation. Only those varieties in the "Good Promise" category could be consistently considered as acceptable to the trade both in the domestic, as well as foreign markets. Data for the named varieties of Butte, Chris, Era, Marquis and Waldron will be an average of each variety for the last three years.

HS 7664

Kernel Characteristics - Questionable to Unsatisfactory. Tendency toward low protein and test weight.

Milling Performance - Satisfactory to Questionable. Tendency toward high flour mineral content at 65% extraction.


Baking Evaluation - Unsatisfactory. Long mix time and low absorption.

General Evaluation - Based on this year's crop results, this selection shows no promise due to low protein and low bake absorption.

HS 74183 (2.0 - 33/10)

Kernel Characteristics - Questionable to Unsatisfactory. Low protein content and a tendency toward a small percentage of large kernels.

* (Average General Evaluation - #Total Deficiencies/Major Deficiencies.)



HS 74183 (Cont'd)

Milling Performance - Satisfactory.

Baking Evaluation - Questionable. May have a tendency toward low absorption and a tough dough.

General Evaluation - Based on this year's crop results, this selection shows <u>little promise</u>, mainly because of its questionable kernel characteristics.

The Prospect - Based on three crop years, this selection shows <u>little promise</u>, due to questionable kernel characteristics.

ID 0167

Kernel Characteristics - Questionable. Low protein content and a tendency toward high percentage of small kernels.

Milling Performance - Satisfactory.

Baking Evaluation - Unsatisfactory. Long mix time and a tendency toward a very strong dough.

General Evaluation - Based on this year's crop results, this selection would show no promise, due to its tendency for a high percentage of small kernels, long mix time and tough dough.

MN 7222 (2.1 - 25/9)

Kernel Characteristics - Questionable to Unsatisfactory. Tendency toward low protein content.

Milling Performance - Satisfactory.


Baking Evaluation - Questionable to Unsatisfactory. Low absorption.

General Evaluation - This selection based on this year's crop would show <u>little promise</u>, because of low protein and bake absorption.

The Prospect - Based on the last three crop years, this selection would show <u>little promise</u> as a new variety, due to its tendency toward low protein and bake absorption.

MN 7357

Kernel Characteristics - Questionable. Tendency toward low protein content.



MN 7357 (Cont'd)

Milling Performance - Satisfactory to Questionable. May have a tendency toward high flour mineral content at 65% extraction.

Baking Evaluation - Questionable. Has tendency toward a tough dough and long mix time.

General Evaluation - Based on this year's crop results, this selection would show some promise, although it has a tendency for minimum protein content and high flour mineral content at 65% extraction.

MN 70170R

Kerne_ Characteristics - Questionable. Tendency toward low 1000 kernel weight and percentage of large kernels.

Milling Performance - Satisfactory.

Baking Evaluation - Questionable to Unsatisfactory. Low absorption.

General Evaluation - Based on this year's crop results, this selection shows <u>little promise</u>, due to its tendency toward poor kernel characteristics and low absorption.

MN 73168 (1.7 - 22/6)


Kernel Characteristics - Questionable. Tendency toward low test weight and protein content.

Milling Performance - Questionable. High mineral content at 65% extraction.

Baking Evaluation - Questionable to Unsatisfactory. Very strong dough and a tendency toward low absorption.

General Evaluation - Based on this year's crop results, this selection shows <u>little promise</u>, because of its tendency toward low test weight, high mineral content at 65% extraction and a very strong dough.

The Prospect - Based on two crop years' results, this selection would show <u>little promise</u>, due to low protein, a tendency toward high mineral content at 65% extraction, low absorption and a very strong dough.



MT 7648 (1.8 - 29/5)

Kernel Characteristics - Questionable. Tendency toward a high percentage of small kernels.

Milling Performance - Questionable to Satisfactory. Has a tendency for high mineral content at 65% extraction.

Baking Evaluation - Questionable to Satisfactory. Long mix time.

General Evaluation - This selection based on this year's crop shows some promise; however, it does have a tendency toward poor kernel characteristics and a long mix time.

The Prospect - Based on two crop years' results, this selection would show <u>little promise</u> as a new variety, due to poor kernel characteristics and erratic baking performance.

MT 7836

Kernel Characteristics - Questionable to Satisfactory. Tendency toward high wheat mineral and low protein.

Milling Performance - Satisfactory to Questionable. May have a tendency toward high mineral content at 65% extraction.

Baking Evaluation - Questionable. Erratic baking performance. Tendency toward long mix time.

General Evaluation - Based on this year's crop results, this selection shows <u>little promise</u>, because of its erratic baking performance.

ND 569 (3.7 - 4/0)


Kernel Characteristics - Questionable to Satisfactory. Tendency toward low protein.

Milling Performance - Satisfactory.

Baking Evaluation - Satisfactory to Questionable. Tendency toward a very strong dough.

General Evaluation - This selection based on this year's crop results would show good promise as a new variety.

The Prospect - Based on two crop years, this selection would show good promise as a new variety.



ND 572

Kernel Characteristics - Satisfactory.

Milling Performance - Questionable to Satisfactory. Tendency toward low flour extraction.

Baking Evaluation - Satisfactory to Questionable. Absorption and crumb grain down slightly.

General Evaluation - This selection based on this year's results would show some promise. It does have a tendency toward low flour extraction.

ND 573

Kernel Characteristics - Satisfactory to Questionable. Slightly lower protein than checks.

Milling Performance - Satisfactory to Questionable. Flour extraction down slightly and a tendency toward high mineral content at 65% extraction.

Baking Evaluation - Satisfactory to Questionable. Tendency toward a very strong dough.

General Evaluation - Based on this year's crop results, this selection would show <u>some promise</u> as a new variety. Kernel characteristics, milling and baking performance are slightly down from the checks.

ND 574

Kernel Characteristics - Satisfactory to Questionable. Test weight slightly less than checks.


Milling Performance - Questionable to Satisfactory. Tendency toward high mineral content at 65% extraction.

Baking Evaluation - Satisfactory to Questionable. Crumb grain slightly poorer than the checks.

General Evaluation - Based on this year's crop results, this selection would show some promise, although it does have a tendency toward high mineral content at 65% extraction.

ND 575

Kernel Characteristics - Satisfactory to Questionable. Tendency toward low test weight and protein content.



ND 575 (Cont'd)

Milling Performance - Questionable to Satisfactory. Tendency toward high mineral content at 65% extraction.

Baking Evaluation - Satisfactory.

General Evaluation - This selection based on this year's data would show some promise; however, it does show a tendency toward high mineral content at 65% extraction.

NK 75S2631

Kernel Characteristics - Questionable to Satisfactory. Tendency toward low protein content.

Milling Performance - Unsatisfactory. High mineral content at 65% extraction.

Baking Evaluation - Questionable to Unsatisfactory. Tendency toward low absorption.

General Evaluation - Based on this year's crop results, this selection would show no promise as a new variety, due to high mineral content at 65% extraction, its tendency toward low protein content and low absorption.

NK 75S2634

Kernel Characteristics - Questionable. Tendency toward low protein content.

Milling Performance - Satisfactory to Questionable. Tendency toward high mineral content at 65% extraction.

Baking Evaluation - Unsatisfactory to Questionable. Low absorption.

General Evaluation - This selection based on this year's crop results would show <u>little promise</u>, due to low protein content and low bake absorption.

RL 4352

Kernel Characteristics - Satisfactory to Questionable. Test weight slightly less than checks.

Milling Performance - Questionable. High mineral content at 65% extraction.



RL 4352 (Cont'd)

Baking Evaluation - Satisfactory to Questionable. Absorption slightly less than checks.

General Evaluation - Based on this year's crop results, this selection would show <u>little promise</u>, due to high mineral content at 65% extraction.

SD 2700

Kernel Characteristics - Satisfactory to Questionable. Tendency toward low protein content.

Milling Performance - Unsatisfactory. High mineral content at 65% extraction and a tendency toward low flour extraction.

Baking Evaluation - Questionable. Long mix time.

General Evaluation - Based on this year's crop results, this selection would show no promise as a new variety, due to high mineral content at 65% extraction and long mix time.

SD 2835

Kernel Characteristics - Satisfactory to Questionable. Tendency toward low protein content.

Milling Performance - Satisfactory to Questionable. Flour extraction slightly less than checks.

Baking Evaluation - Satisfactory to Questionable. Crumb grain slightly less than checks.


General Evaluation - This selection based on this year's crop results would show some promise as a new variety; however, it does show a tendency toward less protein content than the checks.

SD 2868

Kernel Characteristics - Satisfactory to Questionable. Tendency toward a small percentage of large kernels.

Milling Performance - Satisfactory.

Baking Evaluation - Satisfactory to Questionable. Tendency toward a weak dough.



SD 2868 (Cont'd)

General Evaluation - Based on this year's crop results, this selection would show some promise. The baking quality was slightly less than the checks.

SD 2870

Kernel Characteristics - Satisfactory to Questionable. Tendency toward small percentage of large kernels.

Milling Performance - Questionable. Tendency toward high mineral content at 65% extraction and slightly lower flour extraction.

Baking Evaluation - Questionable. Tendency toward a weak dough.

General Evaluation - This selection based on this year's crop data would show <u>little promise</u> as a new variety, due to its questionable milling and baking performance.

WA 6756

Kernel Characteristics - Questionable to Unsatisfactory. Tendency toward low test weight and protein content.

Milling Performance - Questionable to Unsatisfactory. High mineral content at 65% extraction.

Baking Evaluation - Questionable to Unsatisfactory. Tendency toward long mix time and low absorption.

General Evaluation - Based on this crop year's results, this selection would show no promise, because of its tendency toward low protein content, high mineral content at 65% extraction and long mix time.

WA 6758

Kernel Characteristics - Unsatisfactory to Questionable. Small percentage of large kernels and a tendency toward low test weight.

Milling Performance - Satisfactory.

Baking Evaluation - Unsatisfactory. Low absorption.

General Evaluation - Based on this year's crop results, the selection would show no promise as a new variety, due to its small percentage of large kernels and unsatisfactory baking evaluation.



X 6718

Kernel Characteristics - Questionable. Tendency toward low protein content.

Milling Performance - Satisfactory to Questionable. Tendency toward low protein content and test weight.

Baking Evaluation - Unsatisfactory to Questionable. Very strong dough and a tendency toward long mix time.

General Evaluation - Based on this year's crop results, this selection would show <u>little promise</u>, due to long mix time and a tendency toward a very strong dough.


X 6753

Kernel Characteristics - Unsatisfactory to Questionable. Tendency toward a small percentage of large kernels, low kernel weight and low protein content.

Milling Performance - Satisfactory.

Baking Evaluation - Unsatisfactory to Questionable. Tendency toward low bake absorption.

General Evaluation - This selection would show <u>little</u> <u>promise</u> as a new variety based on this year's crop results, due to its tendency toward low protein content and low bake absorption.



1980 UNIFORM REGIONAL HARD RED SPRING WHEAT NURSERY SAMPLES NOT INCLUDED IN THE AREA BLENDS

These samples were milled on the Brabender Quadrumat Jr. Mill, and a 25 gram flour sample was baked. Butte, Chris and Waldron were the standards.

TETONIA, IDAHO SAMPLES

The effect of frost on these samples was evident in the kernel appearance, when looking at the data and in the general evaluation. Test weights in most cases were low and wheat ash showed a tendency to be high. Flour extraction and mineral at 65 percent extraction were severely affected. Flour extraction was very low, and the mineral content at 65 percent extraction was high. The frost had a definite affect on the bake evaluation. All samples showed a weak to dead dough, crumb color was down and crumb grain and loaf volume were severely affected. Data for these samples are given in Table 5.

PULLMAN, WASHINGTON SAMPLES

Several selections from this station showed good promise. However, one must consider that the standards used were comparatively grown, and that the wheat and flour protein levels were considerably lower than desired. This gives a somewhat fictitious general evaluation to the samples. Data for these samples are given in Table 6.

MADISON, WISCONSIN SAMPLES

Several selections from this station also showed good promise. But again, the standards used were comparatively grown and test weight, wheat and flour protein, bake absorption, dough character and loaf volume were severely affected by sprout damage and would not produce an acceptable product. Data for these samples are given in Table 7.

WILLISTON, NORTH DAKOTA SAMPLES

The 1980 milling and baking standard was used as the standard. Coteau showed no promise, due to its poor kernel characteristics. Data for these samples are given in Table 7.



FIELD PLOT NURSERY SAMPLES - 1980 CROP

Thirty-one samples were received from two states and two stations. The data for the individual samples are given in Tables 8 and 9.

NORTH DAKOTA SAMPLES

Twenty-four samples were received from the Williston station. Sixteen of the samples were named varieties which have been released. The data are given in Table 8. Butte, Olaf and Waldron were used as standards.

ARIZONA SAMPLES

Seven samples were received from the Mesa station. Two of the samples were named varieties which have been released. Cajeme 171 was used as the standard. Data for these samples are given in Table 9.



INTERNATIONAL SAWFLY NURSERY SAMPLES - 1980 CROP

Fifty-eight samples were received from two stations in Montana and two stations in North Dakota. Twelve samples were received from each of these stations: Bozeman and Havre, Montana; and Minot and Williston, North Dakota. of these samples were the named varieties - Chris, Fortuna, Lew, Thatcher, Tioqa and Waldron. Six of the samples were the selections MT 783, MT 7620, MT 7810, MT 7819, MT 7926 and ND 580. The Minot, North Dakota station also included MT 810 and SU 28 SIB-1. The Williston station included Butte, Olaf, MT 781 and SU 28's 1107. Bozeman and Havre, Montana stations included MT 781 and SU 281. Sample quality data from the individual stations are given in Tables 10 through 13. Table 14 contains averages of these data. Again, averages and blends may not reflect the range of response of a selection or variety to environmental conditions; therefore, averages of the general evaluation, numbers of total deficiencies, and the number of major deficiencies are given, the same as they were for the Uniform Regional Nursery series. The varieties Fortuna, Thatcher and Tioga grown at each station were used as the standards.

MT 783 (2.4 - 34/6)


Kernel Characteristics - Questionable to Satisfactory. Low percentage of large kernels and tendency toward low kernel weight.

Milling Performance - Satisfactory to Questionable. Tendency toward high mineral content at 65 percent flour extraction.

Baking Evaluation - Questionable to Satisfactory. Long mix time.

General Evaluation - This selection based on this year's crop results would show <u>little promise</u>, due to poor kernel characteristics and long mix time.

* (Average General Evaluation - #Total Deficiencies/#Major Deficiencies.)



MT 783 (Cont'd)

The Prospect - Based on two crop years' results, this selection would show <u>little promise</u> as a new variety, due to poor kernel characteristics and its tendency toward long mixing time.

MT 7620 (1.8 - 45/13)

Kernel Characteristics - Questionable. Low percentage of large kernels and 1000 kernel weight.

Milling Performance - Questionable to Satisfactory. Low flour extraction.

Baking Evaluation - Questionable to Satisfactory. Tendency toward long mix time and poor crumb grain.

General Evaluation - This selection based on this year's crop results would show some promise. Its kernel characteristics are questionable, and the milling performance is down.

The Prospect - Based on two crop years' results, this selection would show <u>little promise</u> as a new variety, due to poor kernel characteristics and milling performance.

MT 7810 (1.4 - 40/21)


Kernel Characteristics - Questionable to Unsatisfactory. Tendency toward low 1000 kernel weight and percentage of large kernels.

Milling Performance - Questionable to Satisfactory. High mineral content at 65 percent flour extraction.

Baking Evaluation - Satisfactory to Questionable. Tendency toward long mixing time.

General Evaluation - This selection based on this year's crop results would show <u>little promise</u>, due to its poor kernel characteristics and milling performance.

The Prospect - This selection based on two crop years' results shows no promise as a new variety, due to low percentage of large kernels, 1000 kernel weight, high mineral content at 65 percent flour extraction and tendency toward long mix time.



MT 7819

Kernel Characteristics - Questionable. Tendency toward low percentage of large kernels.

Milling Performance - Satisfactory to Questionable. Has a tendency for high mineral content at 65 percent flour extraction.

Baking Evaluation - Unsatisfactory. Low bake absorption and a very long mix time.

General Evaluation - This selection would show no promise based on this year's data, due to low bake absorption, long mix time and questionable kernel characteristics.

MT 7926

Kernel Characteristics - Questionable to Unsatisfactory. Tendency toward low percentage of large kernels, 1000 kernel weight and wheat protein.

Milling Performance - Satisfactory.

Baking Evaluation - Questionable. Tendency for low bake absorption and erratic dough characteristics.

General Evaluation - Based on this year's crop results, this selection would show <u>little promise</u>, due to low bake absorption and questionable kernel characteristics.


ND 580

Kernel Characteristics - Satisfactory.

Milling Performance - Satisfactory.

Baking Evaluation - Satisfactory to Questionable. Somewhat erratic results.

General Evaluation - Based on this year's crop results, this selection would show good promise.



INTERNATIONAL NURSERY - 1980 CROP

Sixteen samples were received from St. Paul, Minnesota. Era was used as the standard. Two selections, MN 7086 and Q.T. 4081 showed good promise; however, Q.T. 4081 was 2.2% lower in flour protein than MN 7086. Major faulting factors were low flour extraction, bake absorption, loaf volume and protein content. Data are presented in Table 15.



EXPLANATION OF ABBREVIATIONS LISTED UNDER THE HEADINGS AND THOSE THAT MAY BE LISTED UNDER MINOR AND MAJOR DEFICIENCIES ON COMPUTER PRINTOUT

TW = Test Weight

KW = 1000 Kernel Weight

LG = Large Kernels

MD = Medium Kernels

SM = Small Kernels

WM = Wheat Mineral

WP = Wheat Protein

KERN CHAR = Kernel Characteristics

EX = Flour Extraction

M65 = Mineral at 65% Flour Extraction

FLR. PRO = Flour Protein

PD = Protein Drop Between Flour and Wheat

MLG. CHAR = Milling Characteristics

MLG. PER = Milling Performance

MIX. ABS = Mixograph Absorption

MIX. PAT = Mixograph Pattern Score

BAKE. ABS. = Actual Bake Absorption

BA = Bake Absorption

MIX TIME = Actual Dough Mixing Requirements

MT = Mix Time

DOUGH CHAR = Dough Handling Characteristics

DO = Dough Characteristics

CRUMB COLOR = Example - 100.5

100 = score received for color

.5 = creamy-the characteristic

of that particular loaf.

CRUMB GRAIN = Example - 86.05

86 = score received for crumb

grain.

.05 = open-or characteristic of

that loaf's crumb grain.

LOAF VOL = Actual Loaf Volume

LV = Loaf Volume

BAKE EVAL = Bake Evaluation

GEN. EVAL = General Evaluation



FARE 1 DIMLITY DATA IF UNIFORM REGIONAL MIRSERY DLENDS

44338 DEFICIENCY		\$ 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	E CO C E C C C C C C C C C C C C C C C C	1 4 5 00	# 55 G A	H1 1 1 1 2 1 4 1 1 1 1 1 1 1 1 1 1 1 1 1	KO HON DA KO HON DA
MINDR DEFICIENCY		HAS LG H65	MAN ME ME ME MEN ME MEN MEN MEN MEN MEN ME	WP H55 DO SH NS NS NS NA 30	EX MOS JA HOS	24 52 45 52 45 52 54 55 54 55 55 55 55 55 55 55 55 55 55	LG SW AT LG SW AT SH Nes RT DO Rd LG SW MASS
GEN. EVAL.		ee-ne		-~~-	N4PR-		4N-
VOL. EVAL.		****	***		4 NN NE	DNDN4	****
134°	į	80000 60000 80000	00000 00000	923 970 970	20000 20000	00000	00000
2000		00000 00000 00000 00000 00000 00000 0000	00000	07.05 05.07 07.07 07.07	84000 10000 10000	00000 10000	\$00000 \$0000 \$0000 \$0000 \$0000
0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		00.00	*****	00000		10000	20000
CHAR. C		nnnnr		POWER	nnene		renor
*5		25555	00000	6.0000 6.0000 5.0000		00000 00000 00000	2000-
435.	-	*****		00000 00000	-05.00 -05.00 -05.00	*****	4 P. C.
11.						****	****
157		55.75	N0000	5	20000 20000	22.52	
200		พพคสก	DNEDD	-PF-eN	nmere	040 ^m N	PONEN
FLO. M.G.		Denve Denve	PP4-PP	54.0E			2222
A SKE	-		00000	00000	00000	00000	20000
:: ::3		- 2.5.0 - 2.5.	46-55	12001	-0000 -0000	-0.001	00000
THAR.		~~~~	***	****	~~~~	*****	700r
PRO.				40000	*****	99996	200-0 245-0
# F F F F F F F F F F F F F F F F F F F		00000 00000 00000	-0000	50000	20000	00000	5500 6600
35	_		n	~			-~~~
FEBRESUES STEE			*****	0 0 0 0 N	4 & S & Line S = F & Line	*****	6-P40 6-P40
		-4000	0-0-0 0nn-c	*****	20205	40000	25020
0.	10.0	20000	24.55	30000	80000		242
<u>:</u> -		20000		00000	6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	100000	*****
VARIETY OR	NOBTHE ASTERN	91116 CH115 804 MATOUIS KALDRIN	15 7664 19 1067 19 1222 14 7222	0 C C C C C C C C C C C C C C C C C C C	NO 572 NO 573 NO 573 A 755 2631	NK 755 2636 91 2733 93 2635 50 2835	0 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

C.EAN DOY - SUBJUCT 1 13.770. FOR DICKAGE-FREE 1.4.

1 A MOTE DATA - 2 ASSTRACTORY - 2 ASSTRACTORY - 2 USSTRACTORY - 5 OFFICE OF



FABLE 2 BUALITY DAIA JF WHIFUAM RESIONAL MURSORY BLENDS

APPENDING SELECTERCY	ž	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	** **	8 U X	**************************************	46 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
MINION DEFICIENCY		4 A OF	0±			
2	46 2 3 A	1280	315	# 0 0	Ï	_
BUN .		3375	7401 6511	3 4		52707
, ,	2222	22226	22000	0£ ±3	2010	22233
GEN.	りゅいかり			MN-M-		
BYAL.		04004	engpe	44FM0	****	00000
100A V01.	0000E	20006	*****	*****	*****	Popor
MI PUS GRAIN	000000 00000 00000	00000	7.000 0.000 0.000 0.000 0.000 0.000		00000	000.000
C3000 C3C3	2000		-0000		*****	*****
00064 CHAA.	****	PPwns	m&866	nnnne	nnnnn	ennun
1 tr	###### ###### ########################	647.22 67.22	75.54 5.55 5.55 5.55	845/45/ 808/86/ 808/86/	nocno nrono enter	2007-4 2077-0 20
A CONTRACTOR	****** ****** *****	00000	05-N4 10-55	64000 00000 65000	00000	20000 20000 20000
74	SUBSTITUTE OF		nee-e	ขพพ64	84280	****
750	00425	5000C	800000 800000	00000	00000	20000 20000 20000
Pen.	NNPER	กคพกพ	PEKNO	Panes	POOPL	50 10 10 10 10 10 10 10 10 10 10 10 10 10
CHAN.						
12. 1.0.	24060	2000	2222		N4555	00-44
7242	00000	0000C	*****	00000	700NE	N4-PN 444
7 L	500000 50000 50000	0-00F	F0000	00000	00000	NO 0 0 P
SHAR.	PROCE	****	****	*****	O D449	noon
E COM						
7 Nam	00000	50000		 	20.00	
15 P	20000			****		0-000
10 M	40270	2017 2017 2017 2017 2017 2017 2017 2017		00000	2000	01077
1 !						
6 E		27.000				
, j.ĥ.	2 00000	######################################	200000 200000 200000	00000000000000000000000000000000000000	**************************************	**************************************
VANETY DA	SCUTTE STERM BJTTE CHRIS E14 MARBULS	9001 0 1 7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2000 2000 2000 2000 2000 2000 2000 200	42 872 43 874 43 874 43 874 44 785 44 785	4x 785 2694 4, 4352 5) 2703 5) 2018 5) 2640	SOUND THE SOUND SO

CLEAM ONY - SUBTRACT I .5.730. FOI DOCKAGE-FREE T.W.

1 MIN STREAM ON - STREAM



DJALLIY DAFA 3F UNIFORM REGIONAL NUMBERY SLENDS PABLE 3

NAJOR DEFICIENCY			0 S S S S S S S S S S S S S S S S S S S	* 4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	n 0 1	40 8	40 %	egy ca egy ca egy sign
MINDS DEFICIENCY	* 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		7 7 7 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	No N	SM NF NF DO LEG SM NF DO BD DO BD DO	65 00 100	2 2 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1.6 00 1.6 NC 1.6 S.4 86 S.4 87 UD
GEN.			44004	-~-~~	48000	4 W W T =	08-00	00
BAKE BVAL			0 N 4 N N	04644	NNFFF	*****	04844	
LOAF V3L.	2		4000	00000	20200	95000	00000 00000 00000	00000
CAUMB 641463			00000	000000 000000 000000000000000000000000	00000	00000 00000	10040 6040 10040	00000
0.08				2222	*****	*****	2000 2000 2000	*****
5000			00000	20000			DDD44	
**************************************			2000 v	48.44.0 64.44.0 64.44.0 64.44.0	00000 00000 00000	20000 20000 20000	25.23	20.00 30.00
94KE 405.			70	0045. 0045.	2000	00000 00000 00000	88 - C 4	**************************************
***			N+NN0	****	40.00	84848	40-44	
485.	-		00000	55555	20000	20000	20002	******
MLG. PER.			พพกตก	+n-n+	NEADN	Panad		@ENMN
CHAR.								
FL9.	ger			20000	14111 14111	-0-00		22222
MIN.B 65KEK.	_		00000	00000	*****	2222	00000 00000 00000	4-600
ENT. 65KE	i Der		-4400	-0150	20200 20200 20200	99999	2000 6600 1000 1000 1000 1000 1000 1000	7.00.1
KEPU. CNAR.			N4 0 NN	enene	n04+0	****	ยคพพค	nadea
PRO. C	!		33.75		00000	11050	14882	0-202
22.2	!		00077	90000		20220	00400	20000
LET Name	34		~~~~	22244	2448W	N45~4	00000	n-sina
Egangle stiff	in the last			20000	*****	20000	22500	200
	į		22426	242-0	00000	-80-2		-6-66
95			22022	20.50	9-225	20074	******	22.26.03
<u>:</u> -		CN	00-00 00-00	20008	77=F4 200000 200000	00000	******	-2000
VANTEN CH		WESSEAN RESEGN	BITTE CIRIS FER HARDIS	112 7 65 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	11 70179R 11 70179R 11 70165	N 20 0 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	4K 755 2634 R: A354 S0 2733 S0 2835 50 2835	S) 2676 4 4 6756 8 5756 8 5753 8 5753

CEEM DAY - 5110ACT | -0.700. FOR DECAGE-112 TOW.

1. AND TOWN STATEMENT OF THE STATEMENT OF



TANJE 4 GUAL ITY DATA OF UNIFORM REGIONAL MIRSFRY MLEMOS

MAJOR DEFICIENCY		- S S		NP BA				₩ B
WINDO DEFICIENCY		WF 00 LA SH M65 MH M65 SH M65 0A		148 55 185 ms		SH SH NGS BA HIGS SM MGS		25555 25555 25555 EA
GEN. EVAL.		name				++++		
7. K.				****		*****		
LDAF BANE VOL. EVAL		000000		66444 88800		0000		50 50 50 50 50 50
N 7 6 5 7 6 7 6		000000		23.00 23.00 27.00 7.00 7.00		65.00 65.00 65.00 65.00		00000 00000 00000
בא נמנימה 1. ניונים		200-1-00-1-1-00-1-1-1-1-1-1-1-1-1-1-1-1-		00000		7.000		- 000c
CHANG.		*****		*****		90000		00000
=======================================		DD444		40444		WW WW		00000 00000
BARE A05.	,	50000 50000		0000¢		00000 00000		00000 00000 00000
#5.		***		00000		21 4 23 20 ED		****
485.	,	00000		00000		00000		#00kk #8866 9066
PES.		wene		-		****		Column series
M.0 CH4A.								
FLD.	,	04004		2000		****		50000
A THE R	ı	00000		00000		00000 ULUEU E 6801		00000 00000
===	,	00000		722.2		70-07		00000
KERN. CHAR.		N M4MB		Ne sont		*****		No see con de l'an
Pag.						******		4044 4044
I Z				*****		2007-		Pp 0=0
KERNEP 8185		rcoor				****		*****
NE C		TERRI				****		2004t
E .		24982				2-19t		850NN NEDED
X 4 4 .	PAGES	22022	PAGES	0 F C C C C C C C C C C C C C C C C C C	-	27.5		500-0 600-0
1000 1000 1000 1000 1000 1000 1000 100	OF & AV	CENNE CONTE CONTE	OFA AVE	C664-	VERAGE	500000000000000000000000000000000000000	MGFS	00000
VARIETY OR 971. MD.	SHITTER STEPH ANEA AVERAGES	MITTE CHRIS FRA WARRON ARFA AVFRAGE	NORVIITASTEDN ARFA AVEDAGFS	BILLIF CHOIS ERA WALDODN APFA AVCRACE	WFGIFGN APFA AVERAGES	91111E CIRATS ERA WALDGON AREA AVFRAGE	FROM YEAR AVERAGES	1977 AVERAGE 1978 AVERAGE 1979 AVERAGE 1977-10 AVE

22222222

CLE 44 DOY — SUNTACT I LD. MUL. FOR INCKAGE-FREE T.W.

TO STATUS OF THIS STATES OF THE STATES OF THE STATES OF THE STATES OF THE STATES OF THIS STATES OF TH



						3			:
								2.2	• 'D9
	ž			Ę	Ę	. 1			, ž
	=		5		30	22	2 ¥	400	10 P
	10		0 4 0	#0 XX0	20 AA	2202 2200 2200 2200 2200 2200 2200 220	2 X X 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	2000 2000 2000 2000 2000 2000 2000 200	3r40
	ē							PEXX	1 2 2 2 1 1
	ī								2 4
									4 3 4
									ž :
			3					3,	4 6
	<u>.</u>							Ē	JF51
	Ž		Ì		,			8 8)
			2		ē	2 500	400	8 5	2 22
	6			3 055	3	23322	5 378	## W	Y YEAR
	, ,		25; 3	52225	: ::	2×3	02722	1055	2 0 - Z
	ż								. *****
	W>0		-0-5-						
	JAKE JAKE		@~@nn	****	សមាសមា	****	04444	0040	5 × 5
	ر د ۲		#00h-	00000	8282	00000	00000	S550	S TO S
	55	Ĩ							1 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -
	92		20002	60000	35555	85568		3556	X
	55		00000	25857	20202	22822	00000	20.00	And Care
	04		n=n00	ทุกทุกก		00000	nennn	n n n n	
	555		****	*****	*****		*	25.25	2 70
	5.70		C C O C N	00000		~0500	00000	00	* IXX
		•						•	
	*# **	1.	0.000	22.02	2000	20000	0.000	~ 50 CC	7.44.3 CRE-
	•			•		• • • • • • • • • • • • • • • • • • • •			2 22
	4EV	i i	D+05+	MORKE Neron	*****	20000	Freen	-2005	ž 175
			*****	22007	****	25.65	****	0000	2 - X - X - X - X - X - X - X - X - X -
	***			enene	enenn	40404	****	new<	
			****	nnner	****	20000	rrere		4 -162
	150		50000	20000	90000	25000	20000	-200	75.00
	00	į	@N®N@	E P-0 0 F	0+80-		00000		
		١							¥ 20.55
	125	,							121 121 121 121 121 121 121 121 121 121
FILTER TO THE TABLE TO THE TABL	55	Ĺ	 	N-000	-0046	00-00		::::	00 1 2 - 2
FILE									- 2 - 2 - 2 - 2
	N. N.	-	20000	00000	50000	99.50	00000	9999	1,11,18
	10								7
	25	gar.		0 C me	-66	11.000		-000	20 - 1 × 20
	200		mmenn	****	T-MN-FF	0404D	-	005.0	- 440 X 10 C - 480
	S A A A								0 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -
	100	1	2000	550-4	Pere	E-040	40000	2005	STAN STAN STAN STAN STAN STAN STAN STAN
	i	i				A.S. D.O.S.			F 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
F P P P P P P P P P	1 2 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	u War	27.68.6	-000			2.00.		# # # # # # # # # # # # # # # # # # #
								٠	A THE STATE OF THE
	1	-	# N M N PI	nnenn	mn ner	600000	-5-50	*****	3 001 83577 8086 8168 1169
	23	-	000-0 hhot-	*****	*****	22000		1000	# 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
	KE ST		00	no-ne	22222	sonon	0-L-	*	0 4
	1		-4-1114	NA	•.•.		D		1 2000 H
F F F F F F F F F F	0.7		20000	1000 1000 1000	20002	22000	02000	2000	PANAL PANAL VELY VELY VER
	:	9	ornor					***	PART NAME OF THE PART N
# 1	3	-3/8		20000		60000	-0000	******	A PER DE LA PER
1	•		Q .						15 7 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
L I W OF COURT PRINT CORNER CON CONTROL CONTRO	1 2.		- : ".	m 	0 6	96	NDSE		ZI >ZE O O O OZ
35 # 55031 FACES ALEX 66555 48888 (2xx 799797) 9	-		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	24-07	505	20027 77505 50500	230	2000	
	SAN	i	2000	24252	77:11	95222	5555	* Sxx	



TABLE 6 SUBCIAL UNITRIH HINGERY SAMPLES MOT INCLUOFY IM THE RIZHIS

MAJOR DEFICICATIV									
4 JOB (2 A A	# C		ę	£ ;		
ž									
MINOR BEFICEFUCY))))	; ;			20	, c	
96			22 5	223	0 2			7 5 7 7	٤
			0 0 0 X 0 4 4 I 3 I	212	THOUS	70 I	non I		0 I
GEN. EVAL.			NB-04	··n	••••	eemee		*****	••
BARE EVAL. 32			reenn	7000E	NNFIN E	***	PENP	-	NE
LUAF V3L.	5		7272	100-	2005 2005	10000	31300	00700	**
04.0 04.0 04.0 04.0			000000	00000	25500	20000	50000 50000 50000 50000	00000	86.09
								•	
55 TO			90000	00000			20000		200
0000			-ncor	nnove	****	e emme	neone	enene	••
: b == 1=	Z G		2004 4 20 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	4.05 4.05 5.05 5.05 5.05 5.05 5.05 5.05	88888 80868 88868	**************************************	2.54 2.75 2.75 2.75	*****	6.25
A 15.	ik .		mro-oc 	20120	00000 00000	20025	~*****	CEE.	5.5
110			newns	9P94=	ST S	****	#P###	en in	n >
14	-		88888 8888 8888 8888 8888 8888	20000	20005	20000	25000	20025	63.7
ž.	•		nn	~			-0-00		
2 2 2 2 3									
 			0-000	2-205	202		0.000	0-200	10.2
2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	n-		 	 		70000	00000	00000 40460	0.36
5.			20000 2000 2000 2000 2000 2000 2000 20	88.0 88.0 88.0 80.0 80.0 80.0 80.0 80.0			20000	41-00	54.5
CHAN.			NNEFF	~~~~	≠ NOMP	m wm wm	CHEN SP SHISH	*****	~ ~
	1			2	0000-		-0000		===
ii.	i		50500	50000	# NAWA.	00 NM 6	0-50 h	90000	.33
200			~~	~~	-0-0-	n	wnn	~~	-
1513414E3			1255E	50000 50000	20001	2410 2410	2222	50050	23
E839	-		22440	88764 80-04	40000	90000	*****	F - 0 + 0	50
6. 6.	i		2000 2000 2000 2000 2000 2000 2000 200	94-14 94-14 94-14	D-0000	54 V 55	*******	24.5	33.0
-		1101	AFCOF						63.6
		SHINS				-4006			
VADIETY DR		PULLWAM, WASHINGTON	84716 C-1115 E-24 M494015 SAWYELL	M4L04014 M4MPUM HS 7604 HS 74163	N4 7222 N4 7357 N4 70170 R	4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	11) 575 44 755 2634 44 755 2634 44 4355 50 2733	00000000000000000000000000000000000000	N 3718

.. 6464644



SPECIAL UNIFORM NURSERY SAMPLES NOT INCLUDED IN THE MENDS

WAJOS DEFICIENCY			۵					ž	
32.4			8				20	300 0	
6	-		\$			2	20		
7			ì	355 3	a m	≱ H	3 m m c c	5 W 2	۲
-				ż					٤
2	į			2				m X	ĭ
MINUR DEFICIENCY				75		ä		\$	2.2
8			2 2	£ 5		ž	52	200 J	⊼ tu ≽ à
			94 6 x	72	A.>	2 52	MAT TA	25.420	**
GEN.			88 - 84		40-Ma	4R	n	PA	
BAKE EVAL			*******	880 P N		© 01 - 01 N	N0005	****	N# N
LJAF BAKE WOL. EVAL.	:			2000	-0820	92000		20726	200
CRUMS			90000 90000 90000 90000	00000 00000 00000	2000 000 000 000 000 000 000	0.000	00000 00000 00000	90000 00000 00000	999 999 999 999
CO. 38			00000	0	00000	20000	2000	00000	
Cut 1.					****		-2222	20202	n+n
**************************************	2		467-60	00000 00000	**************************************	04885 78 18	 	8 0 8 0 F	889 880 880
115.			000000 000000 000000	88888 88888 88888 8888 8888 8888 8888 8888	2000E	****** ******	2000 2000 2000 2000 2000 2000 2000 200	******	8000 844
344			80~80 50-05	======	**2-*			***=	one
100	-		00000 00000 00000	200000 200000 200000	00000	88888 80788	*****	2000 2000 2000 2000 2000 2000 2000 200	84.5
. sea.			-N-4F	*~	NN SEN	NP8N4	m 00 n =	+504-	NF-
CHARA									
FLA. Pa3.	-		-0000		-0200	*****		-0006	***
Seren.			00000	0000	20000		00000	00000	000 000
EXT.				62.00	00000 0000 0000 0000 0000	008-00 -008-00 -00-00-00	-04004	00000	500
CHAR.			*****		Nonwe	SI PINO E	ทกพรก	DELNE	~0
980.					0.555	*****	40.00	7.01 0.10 0.10	F-0-0
MEN.			 	01105		*****	, 00 E E	27.45	2 2
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	in a			~~-			~	-~~-~	-~~
1812 - 1818 1818 1818 1818 1818 1818 181	-		50-50 50 50-50 50 50-50 50 50 50 50 50 50 50 50 50 50 50 50 5	27052	20050	25-02	20025	\$000 00000	265
2	-		22525	00000	20275	21005	2022	*****	£ 4
C.			800000 07-000	**************************************	382.43	0-6-66	7,0004	N⊕050= 0000= 0000= 0000= 0000=	33.3 27.0 33.2
<u>;</u>	1730	MISHU	*******	******	************	000000 000000 000000	00000	2000000 200000 200000 200000	96.0 96.0
VARIETY OR SEL. 13.	-	MADISON. WISCOMSIN	BUTTE CHRTS E24 4410UIS 44.00UIS	100 Y 200 Y	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	N 572 NJ 573 NJ 574 NJ 575 NA 755 PAN	NX 735 7634 RL 4357 S) 2700 50 2935 S) 2064	0 2 X X X 4 X 4 X 4 X 4 X 4 X 4 X 4 X 4 X	### 1200 517 NIPPIH 74401A 1700 517 517 517 517 517 517 517 517 517 517

CLEAN DRY - SUMMACE I LD./DD. EUR DOCKAGE-FARE I.W.

1. A MISTALL STATES AND STATES AND



THE COURT CO
NODEN ONDEN DONNE BOSES BEEN NOTED TO THE
WATER CONTROL
TOTAL TOTAL TOTAL STREET STREET
or other cose town grape base
or other cose town grape base

27 | 14 MISTON 0013. | ALIST CONTROL | SALISAGON 1 | SALIS



TABLE 9 21 FIELD PLOT NURSERY SAMPLES

						A - OUESTIONARLE-SATISFACIORY, S - DJESTIONAN,E. B - DJESTIONAN,E. WESTISFACIONT-OLESIJONAR,E. S - UNSATISFACIORY. VERY SOCT. ACT VERY STOONED - VERY SERVICE - VERY CONTROLLE - VERY WEBLIED - STICKLY DEAD. 30 = DEAD. ACT VERY STOONED - VERY STANDARD - VERY WEBLIED - STICKLY DEAD. 30 = DEAD. ACT VERY WEBLIED - VERY CONTROLLE - VERY WEBLIED - VERY CONTROLLE - VERY CONTRO
MAJOR DEFICIFICY			>			
MA JOR DEFICIFICY			34 30	4	0.4	1 OPE
14 JOR			225		EX	16911
1						.
						B.E. S
					£	11344 0.30
» UN U				2	WE KE LO SH WP	V DEA
5			۲ <u>۵</u>	N65 DG	22	CF GRY
, AOX			40 40	5 5	¥0	T SFA WERV XXA-1
Ī			XUX	E.	24	4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
## Z		•		-		* # * * * * * * * * * * * * * * * * * *
BAKE EV4.		04	0=0	c	•0	K KERY K K C C C C C C C C C C C C C C C C C C
FERS. MIG. MIR. MIK. 04KE MIK. DOUGH CRUMB CRUMD LOAF DARE GEN. PPD. (14) PG. A05. AAY. ARS. 114E CR3; CD.OR GRALV V). EVA. BALL MINOR STICENCY A. 37. 27. 37. 27. 37. 27. 37. 37. 37. 37. 37. 37. 37. 37. 37. 3	5	57.0	200	000	790	A = OUESTIONABLE-SATISFACTORY, 5 = DJESTIONABLE-UNSATISFACTORY, 7 = JHSATISFACTORY-OLESTIDAABLE, 5 = UNSATISFACTORY-OLESTIDAABLE, 5 = UNSATISFACTORY-OLESTIDAABLE, 5 = UNSATISFACTORY-OLESTIDAABLE, 7 = UNSATISFACTORY-OLESTICA
22			000	00	67.03	E-UNS
55		é	800	8	00	244. T
2007 2007		3	050	:	**	5 5
200	-	=		-	22	. 1xx
85				_	or	# ## # ## # ##
ž:	H H	3	***		 	F ONA PY CR
8 X E		0.		3	45.	0.1ES
64		•	***	•	۰e	. 5 XXII 3 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
77.		•	***	er.	••	7 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
¥94	East !		5000	6.2.5	6.5.5	2000
FLR. MLG MLO. MIN. MIN. DAKE PRO. CHAN. PET. ADS. PAY. ARS.		^		c	•	2 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
9		_	~	_		Sul A
		3		=	**	VERY SOFT.
==						47 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
720	-		000		90.00	**************************************
E.E.) 	3	200		588.3	25 . KO
700		•	200	e en	~.en	421094-00251100401E. 617075 5 101177 6
		4	- C	.0	12.4 12.4	100 00 00 00 00 00 00 00 00 00 00 00 00
122	-	:				STEE T
170	1 1 1		50		1.37	SOUTH A CENT
196			10 e 6	16	ø#i	A PARALA SALA
15031	De.			;=	\$ E	
E CHE	*	5	000 to	. 5	88	B . 700 E . 70
THE TY UN 1.5. CAT. ESPECE 114: DAY. CHAY. FAT. AVER.	2				33.9 35 59 AL. 2 35 59	CLIAN DIT - SUDGACT LG./OU. FOR DACKGE-FREE 1.4. W. 1927 547 547 547 547 547 547 547 547 547 54
					63.6	STANCE AND
13	14		200		96	ALAME S
80		FONA				THE SER SER SER SER SER SER SER SER SER SE
VARIETY DR		HESA . AR I FONA	P103RAND 171	-6.43		りゅう
1 > 6	1	# (52.0	240	22	2002262 2



FABLE 10 DJALITY BAFA JF INTERNATIONAL SAWFLY MINSERY SAMPLES

								Jt. A.R.
MAJG3 DEFICIENCY			n o n			14 07	ę.	CLEAN DRY - SJOINCE LOADER FOR STEED THE TOWN STE
I			•			,	•	\$
								MILE: 2 5 = 0E1 8 x x x . 36
								EST10N/ EAD. 3C GULA1.
MINDA DEFICIENCY					ī			AY- 0
1 30			3			_	2	ACT 0
9 OM 6			2 22 2 82	,	22	•	2011	4
			× 1-	,		30	-	. UNS RX . 23 00 EX.
LOAF BAKE GEN.				• •	100	n-		* 940
F BAK			8P#6			••	₽ 11-4	7 - 12 - 12 - 12 - 12 - 12 - 12 - 12 - 1
			2000	- :			0007	344 t s
CRUMS GRA14			0000	00.74	9000	86.65	0000	HANCE IN
30,78 12,18				:		•••		2 - 6 G
MAR. C						ne		# # # # # # # # # # # # # # # # # # #
FIME CHAR. COLDR			67.05 67.05 67.05	: :		9.10	0000 0000 0000 0000	COASTE
BAKE ADS.			2000 2000 2000	2 :			0000 0000	10 10 10 10 10 10 10 10 10 10 10 10 10 1
ABS. P41.			4 01 W	n ,	n e sn	-	~~~	TO THE CONTRACT OF THE CONTRAC
ABS.	_		0000			::	5555	141000 141000
FIR. ML6 MLG. ред. СИ49. РЕВ.			n-NE	~ (ดคล	-~	+N+N	A B C C C C C C C C C C C C C C C C C C
CHAG.				- ·				FT. V
77.				- :	nes	12.3	-522	74 006 114 006 14 14 006 14 14 006
MIN. 0	-		1000 1000	0.42	• • • •	 		A MEN
17. 17.	Der .				0.20		50 FF C	20 - 40 A A A A A A A A A A A A A A A A A A
MIN. WHI. KERN. FLR. MIN.S.			nnne		rn eu m		OPNE	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
15.			2000				-5	7 . 4 . 5 . 4
			0000	1.58	100	0.0	9984	7 GE-FRE 9 HORMAL 5 H
15E			NEWN	~	~~		nowe	MANAGE TO SEE SEE SEE SEE SEE SEE SEE SEE SEE SE
Selena Selena	-		00-1	-	964	200	20 - 20 00 - 20	
200	-		N 0 6	9	~ E e	:=2	70nh	1000 100 100 100 100 100 100 100 100 10
1606 KW1.			9555	9	200	20.0	0000	
VARIETY OF T. W. KWI. PERCEPT SIZE	-:0}(1	TANA	2000	\$ 5.3	50.00	62.5	*****	10/2 2 20 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
P0 .01		BIFEMAN. MINIANA	4 0		z -		٤	A
VARIEL SEL.		RIFEMA	CH913 FORTUNA LEE	1 0 GA	HI TOS		110 242	



FIGURE 11 DJALITE DATA JF INTERHADIONAL SAWFLY 4JASSERY SAMPLES

	ICIFHCY												DEN. I AREGJ	
	MAJOR BEFFEIFHEY			٥	ž	201			5		= JHS411SFAC		" Stightly o	
	MINOR DR. ICIENIC					P0 41 00					SOUTHING E-SATISFACTRAY. S . SUZSTINANDE. 6 . B OJESTFONANDE-UNSATERFACTORY. 7 . UNSATINFACTORY-OUESTINANDE. 3 . JUSTINSFACTORY-OF	. 7 = 3[143] T. XIACA 6 = WEAC-[143] E. 9 = WEAV BENY WEACH PO. = SLIGHTLY DIAJ, 30 = 2643. Alany Mas. 4 - Wedy Coffany, Mis Cary, Was. 2 - Out Gray, Mas. 8 - VERY GIAT.	. WXX.30 = 322N. XM4.10 = IRRESULAR. 4X4.31 = SLIGHTLY OPEN.IRREGI	
1	31.8			2	ž	5 2 2 5		<u>L</u>	۲		ACTOR	95	ti 0	
	E 0 H			2	ž5		S.	2	60		11 84	200	×	
	Ī			3	23	35	20	×	3 x 0 3		SH5	2	¥ .	
	EVAL.			n	PNP#			••	mm		:	A X		
	BAKE BVAL			800	NP#N	~~~	Po	~~	1 64 B)		CFORY	VERV	KK.	
	V). EVAL. EVAL.			90.	- 00 0 - 00 0 - 00 0	500	000	200	200		HSATI SFA	AX . 16 .	4,000	
	GRACH			88.69	10000	95.63	50.	98.00	900		MABLE -IN	9 = WE	PJC Diak	
	102			:	***** *****	00.	900		9.50		3116	191.6		
-	MIX. ODUĞU CADMB			2:	2222	22:	ne	2:	100		- C	44-f.		
-	55			ın ı	rnee						9 - 6	300	UL AB.	
		1				5.5	Ī		3.66		251134A	XEAC + 6	V I PRES	
	BAKE ASS.	1		39.3		500	25	90			3 . 30	LI A 3. E-	St 1640t	
	¥.			•	e one	n~ :	°-=	۰.	n er e		FRAY		N ON	
	ABS. PAT.	-		65.3	5000 6000 6000			90			AT ISFAC	-440		
	9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0			~	Penn	P 04	0 P		40 N		AR. E-9	ASIIC, 6 * OLIAGE	PRESULAT. MAX.06	
	HLG C4AR.			-							51.34		PE 30	
1	FLB. HLG. 4.G.			9.0	 	55.	250	2:	40.0		Yeav	PY 464X	65 a 32E4, Iquisulat, Max.06 = 3116Htly topeGulat, xxx.60	
	EX.			9.47			200	~			HABLES VA. 6. B	1		MI 56.
				-	2000 2000 2000 2000 2000 2000	37		•	100		10		×	PRO
					9999	nn.	nne	9	nen en		-3.53	ABLE.	356	3R00
	E N			•	NN	Per	oon	•			51331	C-PL	- 6	•
	MHT. WINY. WERN. FLP.	-		16.9	onec onec	00	000			. T. W.	54 TI SFA	TE PAT	644.33	10M15
	÷;			1.43	 	66	200	1.17		KAGE-FR	Y. 9 E	3	HAP9H.	= 504E
	1			m	~~ ~~	Ne	•		nne	4 00C	100	NOR BE	.63	
	S CO	-		9	*N 4 4 *D 0 6	200	200	90	.00	J. FO	113	E	¥ K	DX 6 S
	100 100 100 100 100 100 100 100 100 100	-		9	:-n	20	0 0 D	2	N - 4	18.781	2 . 3	STAN		TLE P
	ANT. PERESTER	N)		9.52	9656	*	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	26.6	-95	JACT 1	1317	N K		-
	STREET OF TAX NATURE SINCE ALL PASS. CLARK SILVE			31.0	0000	25.		63.6	 	Y - SUB)	SA 16 5" AC	THE TO RESERVE NAMED FOR NUMBERCAL CURVE PARTERN. II & VE I = 0.CXY, 2 = VERY ELASTIC. 3 = ELASTIC. 4 = ELASTIC. 3 =	2066v .	PONISE
	ž .		HAVGE, XONIAN							4H OR	2011	X 10		
	VAR 1ETE 03			5	F33 LUNA LEX PHATCHER T105A	6 AUM		1626	20 19 20 20 20 20 20 20 20 20 20 20 20 20 20					-
	> 5		7	C 19	ENTE	, x	:::	Ŧ	O T D	2	ù'n.	:66	22	6



TABLE 12
BUSTITY DATA D" INTERNATIONAL SAWELY NUMBERF SAMPLES

	44JOR DEFICIENCY							i	¥ 41	76 c#	LG N35	
	7								•	•	•	
	NC 4						8A 47					
	MINOR OFFICIENCY			٥			6			90		
	98 Oc.				4 0 1				9	5	ï	
				F 1.0	2 ×		2 K	2	٥	¥	2	
	EVAL.			••	~	•	- №	• •	-			•
	BARE			7 N	• *	~	o er	~~	•	0	47	~
	LJAF BARE VOL. EVAL.	!		202	226	2	502 202	226	~ ~	208	200	000
	CRUMO			95.03	65.03	60.99	69.07	63.03	06.01	97.10	01.09	40.00
-	CHUNB			138.0	200	0			105.3	101.0		9.10
1	DINGIC C				-		~~	-	-	•	m ~	•
	12 4K.	!		5.19	7.29	9.50	5, 19	6.00	15.00	5:53	200	9.29
	PAKE 494.	-			5.6	65.3	67.0			\$1.9	•	6.1.0
1	***			••			••	~~	_		£c	
-	#0 # P	-					۰.۰	•	-	6		•
-	7.0	!		4.1.0	23	2	63.2	- 5	•	5	Ź	•
	7 × ×			-~	-	٠.	~	~	•	-		~
	15.					-			-	-		
	FLR.	-				_	14.3		_		- 6	
	SEX.			0.33				-	9	0.39	0.33	
	7.5					0.00	59.9		6.2.	62.5	-	25.
	THE P.	7		~	•	• ~	~ •	•	•	•	~	~
	.000			2.5	9	-		-		9.6	::	
					2		•	=	•	1.21	3:	
	107			me		25	P. d		•		7	• •
	£ 23	-		::	2	e ~		6		6	9	200
	BERNEF SIEE	-		=;	:2	2	==	=:	2	=	5	- 5
	0.5			27.2		33.1	32.9			73.3		20.00
			948014	8.0	39.3	. c	93.9				20.0	9
	TABLETY OF 1.300 EEBNEE 316 MIN. PAG. FAR.	!	MINUT. NORTH	61110	LEV	11064	#4L340N 59.5	200	0	AC 102 A	N) 000	010



-

TABLE 13 OUALITY DATA 3º INTERNATIONAL SAWFLY HUNSEDY SAWPLES

WA JOR DEFICIENCE		2.2 2.00	2 + 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -	EA		FERT SOFF. VERY SIBOAT STATUSK, S = OUESTIONABLE, B = OUESTIONABLE - UVSATISFACTORY, T = JUSATISFACTORY OUESTIONABLE, B = UNSATISFACTORY, FERT SOFF. VERY SEAL STATUS OUESTIONABLE, B = UNSATISFACTORY, ENTER THE CONTROL OF THE CONTROL OUESAY, SEAL STATUS OUE OR VERY SEAL STATUS OUESAY, SEAL STATUS OUESAY, SEAL STATUS OUE OR VERY SEAL STATUS OUESAY, SEAL STATUS OUESA
HAS CONGRECAVED CRUIS 10.0 BARE GEN. MININ DE LCIENCY		e e	**************************************	P 00		FERT SOFF. VERY SIDNY: G = OUESITONABLE, B = GUESTIDWARLE-DYSMISFACTORY, 7 = JVSAIISFACTYR-DUESITOWARLE, R = VERY SOFF. VERY SIDNY: G = GLIGHTLY OFFOR. SOFFOR SOFF
41 438		KV LG	43434 CMM2	27273	*	2
35.		esse	non	444N	-	* 4x * 4x * xq
, K		NN648	- unne	NPMP4	~	CARA CARA CARA CARA CARA CARA CARA CARA
OAF BA	ະ	2006	33350 33350 33350	202	101	1 SFACE
CRUMS		900.01 95.00 95.00 95.00	00.00 00.00 00.00 00.00	90.03 90.03 96.03 90.03	60.20	ARLE - U43 AF 9 v W144. 4 v XX4.2
10 P		04450	00000	20000	101.9	49LE.
1000		nner	DDDD		91	* ***
- V-	1212	7.1.0 7.1.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7	86.50 80.50 80.50 80.50 80.50	20000 20000	3.13	OMABLE. CREGOLAG
94 AE 4BS •		********	40000 40000 40000	₩₩₩₩ ₩₩₩₩ ₩₩₩	6.00	OUESTI LE-WEAA HTLY IR
		2000	e è c c c c	ÖĞMÜC		TY. S.
AUS. HIA.						1 - 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4 -
:	:	20000	00000	50000 60000	69.3	8 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -
FL9. MLS. MLG.		N&2-r	N# M B N	NNO	~	
155					-	50000000000000000000000000000000000000
		cense	1 2 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	10055	=	644 50651 64-6-4-11 64-6-4-11
74.58	-		00000 25025		0.44	2
55		005 625.2 625.2 600.0	00000	000000 00000 00000	9.19	25 - X0
WHY AFRN FLR. MIN. B			NEGRE	E WE WA		1004-23 11004-23 11004-1100 10004-1100
940.		mer4e	244.0	20050 20050	13.3	1.5FAC 1.5FAC 1.5FAC 1.5A11C 1.5A11C 1.5A11C
÷ ;		25225	50-55	99299	<u>-</u>	AN CONTRACTOR OF THE CONTRACTO
With		* N. N. N.	NEO-F	4NN-0	~	CTORY CONTRACTORY CAST
- C-1	ner .	00.00	42000	0.400	:	FDR WHITE
A ST		06202	#0n+D	07E68	=	S. 790 NATES
50.0		04.00	050-0	94884	27.0	A - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -
UP COLUMN	HIH OAKUT.	00000 00000	00000	00000	56.0	CREAMONT - SUBJECT L3.790, FOR JOCKAGE-FREE T.W. 1 - VERT EALS ACTORNY 2 - 91415 ACTORN 3 - 5115 CTORV-3/E311744R E. 4 1 - VERT EALS ACTORNY 2 - 91415 ACTORN 3 - 5115 CTORV-3/E31174H E. 4 1 - VERT EALS ACTORN 2 - 91415 ACTORN 3 - 5115 ACTORN 3 - 9140 ACTORN 3 - 915 ACTOR
VARTETY OR 1.4. (WI.		CHAIS FORIUNA Lint HHATCHER 1106A	441.080N 417000 417000 417010	41 7926 845 891 841 891 84 861	50 28.9 1107	CLEAN CONTROL OF CONTR



TABLE 14 DUALITY DATA JF INICANATIONAL SAWFLY JUNTER SAWFES

44331 DEFICIENCY			3 7	C.	VERY SOLTS VERY STATES AND STATEMENT OF A STATEMENT OF THE STATEMENT OF A STATEMENT OF A STATEMENT OF A WASHINGTOOT OF A STATEMENT OF A STATE
MINDS DEFICIENCY		KW LG 36 KW LG 36	KW LG FW SW DO KW SW LG LG 605 MT WP BA DO	L2 5A 30	FERT SOFT. VERT SIRONG. PRINCE OF STIONABLE. A DIESTIONABLE-UNSALISFACTORY. 7 = UNSALISFACTORY-DIESTIONABLE. 6 = UNSALISFACTORY. FERT SOFT. VERT SIRONG. PRINCE-VERT. BY WAY. STANDARY. PERCENTIAL OF STANDARY. STANDARY
HIN. DOUGH FOUND CRUMB LOAF BAKE GEN.		84886 40066 80066	**************************************	199 ts 3	GFACTORY. 7 = US GULL GRAY. MANN. No. MANA. 50 = UPF
CHUMB LOA	22	00000 0000 0000 0000	r nnono 0 0000- 0 0000-	60.00	T104AR E-U45AF 1.
Ar CHAR. EDLUR	Ri	3.7.4 9.00 9.20 9.20 9.20 9.20	r Hembe	5.50 4 133.0	ANDES A DIES
BAKE AGS.	<u>1</u>	NACC SURVE S	-	8.86	PLIABLE-WEAK. ANNAL - VERY CANAL
A PRINCE)	NAMA SUBSE Subse Subs Subse Subse Subse Subse Subse Subse Subse Subse Subse S Subs S	N M++NN	2 63.5	E-S4 VIS* 1103 9180461 PLIASE: 7 B N.S CREMY NAKE 05 B NOSE
TELS. N.C. BEC. 1915. REIN. ACCT.		0000 0000	. 0	19.5	
		5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	. ****** • • • • • • • • • • • • • • • • • • •	61.0 61.0 0.42	12 FEB 1 T A S S S S S S S S S S S S S S S S S S
RIIT. KERN		**************************************	0 0000-	20.01	A SOFT W. SOFT SOFT SOFT SOFT SOFT SOFT SOFT SOFT
1 5 2 1 5 WILL	7,	+8nn	N Nenen	F-	TON DOLLAGE TON DOLLAGE TON GOVERNMENT TON GOVERNMENT TON TON TON TON TON TON TON TON
KAT: KEBN		23.54 33.52 34.52 34.53 35 34.53 34.		35.1 35.1 35.1	1857 1 15.730. 115.730. 116.730. 116.730. 116.730. 116.730. 117.180. 117.180. 117.180. 117.180. 117.180. 117.180. 117.180. 117.180. 117.180.
AASTITUTO 1.4. KEIL CENTED SEE WITH PRODUCTION FILE STREET		VARIETY AVERAGES CLAIS CLAIS CLAIS CASCUS MACHEN MA		7725 61.9 568 59.3	CLEAN DRY SUPJECT 1 19,730, FOR DECAGE-FREE T.W. - NET PAIR SETTING E. B STIFFT. 1107, 2 - 5115FT. 17-75FT. 149, E. 1 - 57 - NORMAL, 2 - NORMAL, 2 - STIFFT. 1 - 50F-NORMAL 4 - 50FT. 5 - 50FT. 1
VAR T		C 44165 FORTING	2000 2000 2000 2000 2000 2000 2000 200	12	



7481E 15 DUALITY DATA OF INTERNATIONAL HURSERY SAMPLES


NAJNY DEFICIFICY			Hr.9		4A 03 LV		:	
MA Jin			5	Ĕ	Ĩ	¥ E	# E	
>			2				ים ני	
MIND? DEFICIENCY			} 8				465 84	
35.1			2 7					_
Ç			4 6 6 6 4 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6			9 4	2040 2040 2040	P) 30
			***	•	oI.	ءَ د	*0 * ô	۵.
E C			~		n-')-«	mufle e	6
47×			c	~	+ 0 ·	.00	****	<
LOAF DAKE VJ. FVAL.	٤		525	==	22	200		198
04.0 04.0 14.0			000	66.69	88.09	90.00	000000	61.19
0.38			200		90.0	200	20000	9.50
CHA4. CO. 38					4 E	000	00000	
AAKE HIR, DOUG! 29U46 AAS. TINE CHAR. CO. JR				7.50		6.00	44548 64548 64566	4.03
AAKE AAS.			-01	59.3	55.5	5.55		63.5
775			- ۳-	•=	25		****	r
	•			50.0	500	0 0 0 0 0 0 0 0 0 0 0 0	-5000	6.1.5
FLR. NLG. HLG. 545. CH44. 054.			-0.	, 4 0	Ne	r ~ ~	10 00 01 P	~
92.							- ~	-
FLR.	100			0 s 0 v	55	22.2	22.55	
MIN. 8 FLR. MLG MLG. MIX. MIX. 41X.	-		5.5.5	0.	9.0		000000	0.52
FILB.			82.9 82.9	20.5	100	00. 00. 00. 00. 00. 00.	200000 200000 200000	37.0
TAN .			in en e	, ww	~~		- 24 - 24	-
WHI. KFRH.	-		200		5.5		0.000	:
MIN.	-		000	-69		200	70/07	
-45	-		200	-=		0 0 0		. 14
200	-		300	200	100	6 0 0 0 0 0	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	:
Egune 315h	-		~*	· ~ 2	~*	•••	***	. #
C			40	222	25.02	22.5	987-10	35.1
::	1966	INE SOTA	989		60.75		00000	
SECURE ON P.S. MET. ESCUELGES SER WITH PRO. CANGE FATE.		ST. PAUL. MIN	A DU CHRATE S	A4ZA E24 JUNCO m5**	KLELN CHAMAC	PAYON 1761	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	U* 262

CLEMN DRY - SUUINACT I LO./BULY FOR DOCKACE-FREE 7.W.

14. URA SATISFACTORY, 2 MAINTING AND CONTROLLED AND CONTROLLED AND CONTROLLED AND AND CONTROLLED AND



REFERENCE MIXOGRAMS HARD RED SPRING WHEAT



U.S.D.A. SPRING WHEAT QUALITY LABORATORY
FARGO, NORTH DAKOTA





